


NASA TECHNICAL NOTE



NASA TN D-4162

2.1

LOAN COPY: RETURN TO  
AFWL (WLIL-2)  
KIRTLAND AFB, N MEX



NASA TN D-4162

REACTION RATES OF CARBON MONOXIDE  
WITH HYDROXYL RADICALS  
AND OXYGEN ATOMS

*by Edgar L. Wong, Andrew E. Potter, Jr., and Frank E. Belles*  
*Lewis Research Center*  
*Cleveland, Ohio*



0130707

NASA TN D-4162

REACTION RATES OF CARBON MONOXIDE WITH HYDROXYL  
RADICALS AND OXYGEN ATOMS

By Edgar L. Wong, Andrew E. Potter, Jr., and Frank E. Belles

Lewis Research Center  
Cleveland, Ohio

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION

---

For sale by the Clearinghouse for Federal Scientific and Technical Information  
Springfield, Virginia 22151 - CFSTI price \$3.00

# REACTION RATES OF CARBON MONOXIDE WITH HYDROXYL RADICALS AND OXYGEN ATOMS

by Edgar L. Wong, Andrew E. Potter, Jr., and Frank E. Belles

Lewis Research Center

## SUMMARY

A mass spectrometer stirred reactor technique was used to measure the rates of reaction of carbon monoxide with hydroxyl radicals and with oxygen atoms ( $\text{CO} + \text{OH} \rightarrow \text{CO}_2 + \text{H}$  and  $\text{CO} + \text{O} + \text{M} \rightarrow \text{CO}_2 + \text{M}$ ).

For the  $\text{CO} + \text{OH} \rightarrow \text{CO}_2 + \text{H}$  reaction in the temperature range of  $310^\circ$  to  $611^\circ \text{K}$ , the rate constant based on  $\text{CO}_2$  production was  $4.2 \pm 0.8 \times 10^{11} \exp(-1200 \pm 300/RT)$  cubic centimeters per mole per second, where  $R$  is the gas constant and  $T$  is the absolute temperature. The rate near room temperature is lower than other values reported recently. The activation energy of 1200 calories per mole (5000 J/mole) is in good agreement with recent estimates.

For the  $\text{CO} + \text{O} + \text{M} \rightarrow \text{CO}_2 + \text{M}$  reaction, an upper limit value for its rate constant based on  $\text{CO}_2$  production was estimated to be  $4 \times 10^{13}$  cubic centimeters squared per mole squared per second at  $500^\circ \text{K}$ .

Since the  $\text{CO} + \text{OH}$  reaction is considerably faster than the  $\text{CO} + \text{O} + \text{M}$  reaction, trace impurities giving rise to  $\text{OH}$  interfere with  $\text{CO} + \text{O} + \text{M}$  rate measurements, and may be responsible for the large differences in rate constants reported by various investigators for this reaction.

## INTRODUCTION

The oxidation of carbon monoxide to carbon dioxide is an important process in combustion and in the photochemistry of  $\text{CO}_2$ . This oxidation process is catalyzed by traces of hydrogen or hydrogen-containing compounds. The effect is observed in combustion of very dry  $\text{CO-O}_2$  mixtures which are difficult to ignite and burn slowly, whereas moist mixtures ignite easily and burn rapidly (ref. 1). In the photochemistry of  $\text{CO}_2$ , the photodecomposition of  $\text{CO}_2$  to  $\text{CO}$  and  $\text{O}$  is efficient for dry  $\text{CO}_2$ , but inefficient when traces

of water are present (ref. 2). Evidently, the reverse reaction of CO and O to CO<sub>2</sub> is catalyzed by moisture.

In the absence of hydrogen, the elementary oxidation step must be  $\text{CO} + \text{O} + \text{M} \rightarrow \text{CO}_2 + \text{M}$ . From theoretical principles, it is expected that the production of ground state CO<sub>2</sub> by this reaction will be slow, since it requires an electron spin change. The catalytic effect of hydrogen and hydrogen-containing compounds on the oxidation could be due to the reaction  $\text{CO} + \text{OH} \rightarrow \text{CO}_2 + \text{H}$ , where OH has been produced by oxidation of hydrogen compounds. After this reaction, OH could be regenerated by reaction of the atomic hydrogen with atomic oxygen. In contrast to the CO + O reaction, the CO + OH reaction is rapid because spin is conserved. To understand fully the catalytic effect of hydrogen on CO oxidation, accurate rates for both the CO + O and CO + OH reactions are required.

The published results on the rates of CO oxidation by O and by OH differ. No one has measured both reaction rates by the same technique to get reliable relative values of the two. For the CO + O reaction, various workers report results which are in poor agreement (refs. 3 and 4). For the CO + OH reaction, Ashmore (ref. 5) gives the most recent activation energy estimates of 0 to 3 kilocalories per mole (0 to 12.5 kJ/mole) (refs. 6 to 8), while older measurements gave values of 6 to 10 kilocalories per mole (25 to 42 kJ/mole) (refs. 9 to 13). Recently, Dixon-Lewis (ref. 14) using an electron spin resonance technique, reported an OH + CO rate constant value  $k_{10}$  of  $1.15 \times 10^{11}$  cubic centimeters per mole per second at 300° K. Herron (ref. 15), using a mass spectrometric technique, reported a  $k_{10}$  value of  $5.1 \times 10^{10}$  cubic centimeters per mole per second for the same temperature. These two  $k_{10}$  values are actually in much better agreement since the values of the rate constants  $k_2$  for the  $\text{OH} + \text{OH} \rightarrow \text{H}_2\text{O} + \text{O}$  reaction used to derive  $k_{10}$  were not the same (private communication from W. E. Wilson, Jr., of Johns Hopkins Univ.). If Herron had used a  $k_2$  value of  $1.55 \times 10^{12}$  cubic centimeters per mole per second (ref. 14) instead of  $0.75 \times 10^{12}$  cubic centimeters per mole per second (ref. 16) his  $k_{10}$  value would be  $8.6 \times 10^{10}$  cubic centimeters per mole per second. These values are much higher than previous published results.

The purpose of this report is to provide better rate constants and activation energies for these two reactions. Using the stirred reactor-mass spectrometric technique previously employed by the authors (ref. 17) to study reactions of atomic oxygen, measurements were made at temperatures ranging from 310° to 611° K.

## EXPERIMENT AND PROCEDURES

### Apparatus

The 300-cubic-centimeter stirred reactor and its connection to the time-of-flight mass spectrometer are shown in figure 1. A more detailed description is presented in




Figure 1. - Stirred reactor and connection to mass spectrometer.

reference 17. The 300-cubic-centimeter reactor was used to study the  $\text{CO} + \text{OH}$  reaction. A larger, 2055-cubic-centimeter, reactor was used for the slow  $\text{CO} + \text{O}$  reaction. Before either reactor was connected to the mass spectrometer, its inside surfaces were wetted with an aqueous 10-percent phosphoric acid solution. The reactor was then dried before being connected to the mass spectrometer to make reaction kinetic measurements. A rhenium filament was used in the ion source to make mass spectrometer measurements because  $\text{CO}$  could not be monitored when tungsten was the filament wire.

## Gases

Carbon monoxide, with a stated minimum purity of 99.5 percent, was sent through a liquid-nitrogen trap before being used. 2.16-Percent  $\text{H}_2$  in argon mixture and a 1.46-percent  $\text{O}_2$  in Ar mixture were premixed tanked gas mixtures. For both gas mixtures, mass spectrometer analysis showed only trace impurities. Analysis of the 1.46-percent  $\text{O}_2$  in Ar mixture showed the presence of 9 ppm  $\text{H}_2\text{O}$ .

Carbon dioxide, with a stated 99.99-percent purity, was used without further purification to calibrate the mass spectrometer for  $\text{CO}_2$ . Hydrogen gas, with a stated purity of 99.9 percent, was also used without further purification.

Nitrogen dioxide ( $\text{NO}_2$ ) gas for O-atom and H-atom titrations was prepared as needed by adding an excess of  $\text{O}_2$  to purified  $\text{NO}$ . The  $\text{NO}_2$  was then condensed out at dry-ice temperature and the excess  $\text{O}_2$  removed by vacuum pumping.

Oxygen gas, with a stated purity of 99.5 percent, was sent through two liquid  $\text{N}_2$  traps before being subjected to a microwave discharge.

## Production of OH Radical for CO + OH Reaction

The hydroxyl radical was produced by the  $\text{H} + \text{NO}_2$  reaction described in reference 18. Hydrogen atoms were produced by subjecting a 2.16-percent mixture of  $\text{H}_2$  in Ar to a microwave discharge. This gas titration reaction was monitored by the mass




Figure 2. - Mass spectrometer monitoring of endpoint for  $\text{H} + \text{NO}_2$  titration.

spectrometer. Figure 2 shows a typical gas titration result. End points can be located by monitoring the  $\text{O}_2$ ,  $\text{H}_2\text{O}$ , or H-atom peak. Our mass spectrometer results are consistent with previous findings (refs. 14, 18, and 19) that the reaction scheme for the  $\text{H} + \text{NO}_2$  reaction at near room temperature is as follows:



Ternary or wall recombination involving these species is too slow to be of importance (refs. 14 and 16). No attempt was made to monitor OH due to its rapid decay (ref. 16). For our experimental condition (2.16-percent  $\text{H}_2$ -Ar, about 35-percent  $\text{H}_2$  dissociated, and a temperature range of  $310^\circ$  to  $611^\circ$  K), the following reaction must be added to the above reaction scheme:



Reaction (4) is relatively unimportant near room temperature but becomes increasingly more important as the temperature is raised.

### Production of Oxygen Atoms for $\text{CO} + \text{O}$ Reaction

In most cases, O atoms for the  $\text{CO} + \text{O}$  reaction were produced by subjecting a 1.46-percent mixture of  $\text{O}_2$  in Ar to a microwave discharge. Some runs were also made with O produced by the  $\text{N} + \text{NO}$  titration reaction and by subjecting 100 percent  $\text{O}_2$  to a microwave discharge.

Mass spectrometer sensitivity to O was obtained by the usual  $\text{NO}_2$  gas titration (ref. 17).

### Calculation of Rate Constants

CO + OH reaction. - The bimolecular rate constant  $k_{10}$  for the  $\text{CO} + \text{OH} \rightarrow \text{CO}_2 + \text{H}$  reaction occurring in a stirred reactor is

$$k_{10} = \frac{-\Delta[\text{OH}]}{\Delta t [\text{CO}]_t [\text{OH}]_t} \quad (5)$$

where  $-\Delta[\text{OH}]$  is the disappearance of OH due to reaction with CO,  $\Delta t$  is residence time,  $[\text{CO}]_t$  is concentration of CO after reaction with OH, and  $[\text{OH}]_t$  is concentration of OH after reaction with CO.

Since OH was not monitored,  $-\Delta[\text{OH}]$  was replaced with its equivalent  $+\Delta[\text{CO}_2]$ , the amount of reaction product produced. The following experimental procedure was used for the rate constant measurement:  $\text{NO}_2$  was metered into the reactor through inlet 2 (fig. 1) to titrate H atoms entering the reactor through inlet 1. Then the  $\text{NO}_2$  flowrate was adjusted to a point just short of the end point, so that no  $\text{NO}_2$  peak could be detected

mass spectrometrically, but only a slight excess of H atoms were present. While this measured NO<sub>2</sub> flow was maintained, CO was then added to the reactor through the same tube. On addition of CO to the reactor, the mass spectrometer indicated that CO<sub>2</sub> and H were produced, and at the same time the H<sub>2</sub>O and O<sub>2</sub> concentrations decreased. This observation indicated that the CO + OH reaction is fast enough to compete for the available OH and that the following reactions are occurring in the reactor:



From the above reaction scheme, [OH]<sub>t</sub> for the stirred reactor in the presence of CO can be calculated. The [OH]<sub>t</sub> calculation makes use of an expression which was derived as follows:

$$\text{Total OH decay} = -\frac{d[\text{OH}]}{dt} = 2k_2[\text{OH}]_t^2 + k_3[\text{OH}]_t[\text{O}]_t + k_4[\text{OH}]_t[\text{H}_2]_t + k_{10}[\text{OH}]_t[\text{CO}]_t \quad (7)$$

By assuming steady-state condition for the oxygen atom, equation (7) is equal to

$$3k_2[\text{OH}]_t^2 + k_4[\text{OH}]_t[\text{H}_2]_t + k_{10}[\text{OH}]_t[\text{CO}]_t$$

For a stirred reactor,

$$\left. \begin{aligned} -\frac{\Delta[\text{OH}]}{\Delta t} &= 3k_2[\text{OH}]_t^2 + k_4[\text{OH}]_t[\text{H}_2]_t + \frac{+\Delta[\text{CO}_2]}{\Delta t} \\ -\left(\frac{[\text{OH}]_t - [\text{OH}]_0}{\Delta t}\right) &= 3k_2[\text{OH}]_t^2 + k_4[\text{OH}]_t[\text{H}_2]_t + \frac{+\Delta[\text{CO}_2]}{\Delta t} \end{aligned} \right\} \quad (8)$$



Equation (8) can be solved for  $[\text{OH}]_t$  such that

$$[\text{OH}]_t = \frac{-\left\{ (k_4[\text{H}_2]_t \Delta t) + 1 \right\} + \sqrt{\left\{ (k_4[\text{H}_2]_t \Delta t) + 1 \right\}^2 + 12k_2 \Delta t ([\text{OH}]_0 - [\text{CO}_2]_t)}}{6k_2 \Delta t} \quad (9)$$

where  $[\text{CO}_2]_t$  is the concentration of  $\text{CO}_2$  produced in reaction (6) and is equal to  $+\Delta[\text{CO}_2]$ ,  $k_2$  and  $k_4$  values are given in references 14 and 16,  $[\text{OH}]_0$  is the total initial OH concentration and is equal to  $[\text{NO}_2]_0$ , and  $\Delta t$  and  $\text{H}_2$  are known.

The rate constant  $k_{10, \text{CO}_2}$  for reaction (6) based on  $\text{CO}_2$  production can now be evaluated by the equation

$$k_{10, \text{CO}_2} = \frac{+\Delta[\text{CO}_2]}{\Delta t [\text{CO}]_t [\text{OH}]_t} \quad (10)$$

An alternate method to calculate  $k_{10}$  is based on the observation that each time more CO is added, the increase in  $\text{CO}_2$  production is accompanied by a simultaneous decrease in water production, designated as  $-\Delta[\text{H}_2\text{O}]$ . This second phenomenon must be due to the competition between reactions (2), (4), and (6) to consume the available OH and is a comparison of the rates of reactions (2) and (4) to reaction (6). This relation can be expressed by the equation

$$\frac{k_{10}[\text{CO}]_t [\text{OH}]_t}{k_2[\text{OH}]_t^2 + k_4[\text{OH}]_t [\text{H}_2]_t} = \frac{[\text{CO}_2]_t}{[\text{H}_2\text{O}]_t} \quad (11)$$

Equation (11) can be rewritten using the term  $R$ , which is the ratio of  $k_4[\text{H}_2]_t$  to  $k_2[\text{OH}]_t$ .

$$\frac{k_{10}[\text{CO}]_t}{k_2[\text{OH}]_t(1 + R)} = \frac{[\text{CO}_2]_t}{[\text{H}_2\text{O}]_t} \quad (12)$$

The term  $[\text{CO}_2]_t$  can be expressed in terms of the  $-\Delta[\text{H}_2\text{O}]$  since, for each mole of  $\text{H}_2\text{O}$  removed,  $B$  moles of  $\text{CO}_2$  can be produced, where  $B$  is a function of  $R$ . For the case

where  $R$  is equal to 0,  $B = 2$ , and where  $R$  is  $\infty$ ,  $B = 1$ . Equation (12) can now be rewritten as

$$\frac{k_{10}[\text{CO}]_t}{k_2[\text{OH}]_t(1 + R)} = \frac{-B\Delta[\text{H}_2\text{O}]}{[\text{H}_2\text{O}]_t} \quad (13)$$

and an alternate rate constant for the  $\text{CO} + \text{OH}$  reaction designated as  $k_{10, \text{H}_2\text{O}}$  can also be calculated since  $-\Delta[\text{H}_2\text{O}]/[\text{H}_2\text{O}]_t$  was measured mass spectrometrically and all other terms are known or can be evaluated.

CO + O Reaction. - The termolecular rate constant for the reaction  $\text{CO} + \text{O} + \text{M} \rightarrow \text{CO}_2 + \text{M}$  occurring in the stirred reactor is

$$k_{14} = \frac{-\Delta[\text{O}]}{\Delta t [\text{CO}]_t [\text{O}]_t [\text{M}]} \quad (14)$$

where  $-\Delta[\text{O}]$  is the decay of O atoms,  $[\text{O}]_t$  is the O concentration in the reactor after reaction with CO, and  $[\text{M}]$  is the total gas concentration in moles per cubic centimeter.

This reaction is so slow that the O decay was less than experimental error. However, the production of  $\text{CO}_2$  could be measured, due to the lack of  $\text{CO}_2$  background ion current. Consequently, the rate constant  $k_{14}$  was estimated by substituting  $+\Delta[\text{CO}_2]$  for  $-\Delta[\text{O}]$  in equation (14).

## RESULTS AND DISCUSSION

### Rate Constants for CO + OH Reaction

When CO and  $\text{NO}_2$  are added through a common tube to H in a reactor as described in the previous section, significant amounts of  $\text{CO}_2$  are produced and can be measured mass spectrometrically. The  $\text{CO}_2$  production for increasing amounts of added CO, expressed as the mole fraction  $F_{\text{CO}}/F_t$ , where  $F_{\text{CO}}$  and  $F_t$  are CO and total gas flow rates, respectively, at a temperature of  $310^\circ \text{K}$  is shown in figure 3(a). The carbon dioxide yield falls off as  $F_{\text{CO}}/F_t$  is increased, and the  $\text{CO}_2$  production expressed in cubic centimeters per second (NTP) is independent of temperature (table I). Occurring simultaneously with  $\text{CO}_2$  production is the decrease in the amount of water due to the competition of reaction (6) with reactions (2) and (4). This effect was also measured mass spectrometrically, and the data for  $310^\circ \text{K}$  is shown in figure 3(b) as a plot of the fraction of  $\text{H}_2\text{O}$  removed

TABLE I. - RATE MEASUREMENTS DATA FOR CARBON MONOXIDE WITH HYDROXYL RADICAL REACTION

[Dilute hydrogen-argon flow (NTP), 1.50 cm<sup>3</sup>/sec.]

Run	Added carbon monoxide flow (NTP) <sup>a</sup> , (F <sub>CO</sub> ) <sub>t</sub> , cm <sup>3</sup> /sec	Mole fraction of added carbon monoxide, F <sub>CO</sub> /F <sub>t</sub>	Temper- ature, °K	Added nitrogen dioxide flow (NTP) <sup>a</sup> , (F <sub>NO<sub>2</sub></sub> ) <sub>t</sub> , cm <sup>3</sup> /sec	Pressure, P		Residence time, Δt, sec	Reaction product CO <sub>2</sub> flow (NTP) <sup>a</sup> , (F <sub>ΔCO<sub>2</sub></sub> ) <sub>t</sub> , cm <sup>3</sup> /sec	Calculated flow of OH (NTP) <sup>a</sup> , (F <sub>OH</sub> ) <sub>t</sub> , cm <sup>3</sup> /sec	Fraction of H <sub>2</sub> O removed, -Δ[H <sub>2</sub> O] [H <sub>2</sub> O] <sub>0</sub>	Moles of water removed per mole of carbon dioxide produced, B (b)	Rate constant, k <sub>10</sub> , CO <sub>2</sub> , cm <sup>3</sup> mole <sup>-1</sup> sec <sup>-1</sup>	Rate constant, k <sub>10</sub> , H <sub>2</sub> O, cm <sup>3</sup> mole <sup>-1</sup> sec <sup>-1</sup>
					mm	N/m <sup>2</sup>							
1a	0.0289	0.0186	310	0.0240	0.70	93	0.173	0.0051	0.93×10 <sup>-3</sup>	0.24	1.93	5.7×10 <sup>10</sup>	4.3×10 <sup>10</sup>
b	.0609	.0385			.70	93	.169	.0084	.86	.36	1.93	4.9	3.2
c	.0993	.0613			.71	94	.167	.0107	.80	.43	1.92	4.0	2.4
d	.1520	.0910			.72	96	.165	.0132	.73	.49	1.92	3.5	1.8
2a	.0238	.0154	311	.0245	.70	93	.172	.0040	.98	.23	1.93	5.2	5.1
b	.0477	.0302			.70	93	.169	.0068	.92	.40	1.93	4.7	5.2
c	.0765	.0475			.71	94	.168	.0095	.85	.49	1.92	4.4	4.2
d	.1226	.0743			.72	96	.166	.0112	.81	.55	1.92	3.3	3.1
e	.1737	.1022			.73	97	.163	.0129	.76	.62	1.92	2.9	2.7
3a	.0232	.0151	444	.0215	.70	93	.121	.0037	.94	.20	1.63	10.2	8.2
b	.0486	.0310	443		.70	93	.120	.0068	.84	.30	1.61	10.0	6.2
c	.0708	.0445	442		.71	94	----	.0090	.77	.44	1.58	9.9	7.4
d	.1066	.0658	441		.71	94	.118	.0113	.70	.53	1.56	9.0	6.2
e	.1475	.0868	---		.73	97	.115	.0134	.61	.58	1.52	9.1	5.0
4a	.0307	.0198	438		.70	93	.122	.0045	.91	.26	1.64	9.4	8.3
b	.0409	.0262	437		.70	93	----	.0064	.86	.39	1.63	10.8	10.8
c	.0698	.0439	436		.71	94	.121	.0086	.79	.47	1.61	9.0	8.1
d	.0936	.0581	436		.71	94	.120	.0105	.73	.59	1.58	8.9	9.1
e	.1315	.0791	436		.72	96	.118	.0119	.68	.60	1.56	7.7	6.2
5a	.0193	.0125	606	.0244	.70	93	.089	.0028	1.07	.09	1.41	14.6	8.0
b	.0403	.0258	607		.70	93	.088	.0052	1.00	.14	1.38	14.2	5.9
c	.0613	.0388	608		.71	94	----	.0074	.93	.24	1.36	14.4	7.6
d	.0993	.0613	---		.71	94	.085	.0104	.83	.40	1.33	14.2	9.2
e	.1280	.0776	610		.72	96	.085	.0122	.76	.49	1.28	14.3	9.2
f	.1727	.1022	---		.72	96	.083	.0140	.70	.58	1.28	13.6	9.9
h	.0334	.0215	610	.0244	.70	93	.088	.0048	1.02	.21	1.39	15.0	12.0
6a	.0559	.0354	611		.70	93	.086	.0074	.95	.31	1.36	16.1	11.7
b	.0829	.0518			.70	93	.085	.0095	.87	.42	1.34	15.2	12.1
c	.1159	.0707			.72	96	.085	.0113	.80	.54	1.32	13.8	12.8
d	.1552	.0924			.72	96	.083	.0133	.73	.61	1.29	13.7	12.9

<sup>a</sup>Normal temperature (300° K) and pressure (750 mm) (100 000 newtons/meter<sup>2</sup>).<sup>b</sup>B is a function of the ratio of [k<sub>4</sub> H<sub>2</sub>] and [k<sub>2</sub> OH].




Figure 3. - Fraction of CO + OH reaction at 310° K.

$-\Delta[\text{H}_2\text{O}]/[\text{H}_2\text{O}]_0$  against  $F_{\text{CO}}/F_t$  where  $-\Delta[\text{H}_2\text{O}]$  is the moles of water removed after addition of CO, and  $[\text{H}_2\text{O}]_0$  is the initial amount of  $\text{H}_2\text{O}$  present before addition of CO. Although the scatter for this water data is considerable, a similar trend, as for the  $\text{CO}_2$  production, was noted. Both sets of data are listed in table I.

The data shown in figure 3 can now be used in equations (10) and (13) to evaluate rate constants  $k_{10, \text{CO}_2}$  and  $k_{10, \text{H}_2\text{O}}$ . The results of these calculations are tabulated in table I and shown in figure 4 as plots of rate constant against  $F_{\text{CO}}/F_t$ . In figure 4(a), the  $k_{10, \text{CO}_2}$  trend with  $F_{\text{CO}}/F_t$  at three different temperatures is fairly well established. The  $k_{10, \text{H}_2\text{O}}$  trends with  $F_{\text{CO}}/F_t$  shown in figure 4(b) can only be determined for the two lower temperatures because the  $k_{10, \text{H}_2\text{O}}$  values at the highest temperature scatter considerably.

These  $k_{10}$  rate constant trends with  $F_{\text{CO}}/F_t$  may be explained by the fact that both  $+\Delta[\text{CO}_2]$  and  $-\Delta[\text{H}_2\text{O}]/[\text{H}_2\text{O}]_0$  tend to level out as  $F_{\text{CO}}/F_t$  is increased and the lack of available OH becomes a limiting factor.

The  $k_{10, \text{CO}_2}$  and  $k_{10, \text{H}_2\text{O}}$  values for the various temperatures were obtained by drawing least squares lines through the appropriate data shown in figure 4. Rate constant values independent of  $F_{\text{CO}}/F_t$  can thus be determined and are tabulated as follows:

Temperature, T, °K	Rate constant, $\text{cm}^3 \text{mole}^{-1} \text{sec}^{-1}$	
	$k_{10, \text{CO}_2}$	$k_{10, \text{H}_2\text{O}}$
~310	$5.8 \times 10^{10}$	$5.3 \times 10^{10}$
~440	10.6	9.6
~610	15.3	-----




Figure 4. - Rate constants variations with added CO.

The  $k_{10, \text{CO}_2}$  values for the three temperatures were estimated to have a precision of at least  $\pm 10$  percent, because all the  $k_{10, \text{CO}_2}$  data (fig. 4(a)) deviated  $\pm 10$  percent or less from their respective least squares lines. The two  $k_{10, \text{H}_2\text{O}}$  values had much poorer precision of about  $\pm 20$  percent. There is good agreement in the  $k_{10}$  values for the two lower temperatures.




Figure 5. - Comparisons of rate constants for the CO + OH reaction.

Using the  $k_{10, \text{CO}_2}$  values tabulated above, an Arrhenius plot of  $k_{10, \text{CO}_2}$  against the reciprocal of the absolute temperature is made (fig. 5), and the rate constant  $k_{10}$  for the CO + OH reaction is equal to  $(4.2 \pm 0.8) \times 10^{11} \exp(-1200 \pm 300/RT)$  cubic centimeters per mole per second, where  $R$  is the gas constant and  $T$  is the absolute temperature. We estimate that our preexponential term has a  $\pm 20$  percent error and that  $\Delta E$ , the activation energy, has about a  $\pm 25$  percent error.

### Comparisons of Rate Constant Measurements for CO + OH Reaction

Rate constant values for the CO + OH reaction obtained by various investigators are also shown in figure 5. The low-temperature data were all from flow discharge experiments using tube reactors, and high-temperature data were mostly from flame studies. The recent low-temperature measurements of references 14 and 15 and the results pre-

sented herein suggest that Avramenko's results are questionable due to the use of a water discharge as an OH source (ref. 16). At 300° K our  $k_{10}$  value of  $5.4 \times 10^{10}$  cubic centimeters per mole per second are lower than Herron's recalculated value of  $8.6 \times 10^{10}$  (ref. 15) and Dixon-Lewis and Westenberg's value of  $1.5 \times 10^{11}$  (ref. 14). The low  $k_{10}$  value reported herein may be due to incomplete mixing in the reactor because of the very rapid reactions. It is interesting to assume that, in the reactor, very little mixing is occurring, and treat the data for 300° K using an unstirred reactor concept. To make this calculation, an integrated expression for a tube reactor (ref. 15) is used.

$$k_{10} = \frac{+\Delta[\text{CO}_2]}{[\text{CO}] \int_0^t [\text{OH}] dt} \quad (15)$$

Equation (15) was integrated using an expression for the OH decay which omitted the additional OH loss due to the  $\text{CO} + \text{OH}$  reaction.

$$k_{10} \approx \frac{1.3 \Delta[\text{CO}_2]}{[\text{CO}]} \frac{k_2}{\text{Log}(3k_2[\text{OH}]_0 \Delta t + 1)} \quad (16)$$

Using the above equation, a  $k_{10}$  value of  $\approx 1.3 \times 10^{11}$  cubic centimeters per mole per second was obtained. This value would be higher if a more exact expression for the OH decay had been used in equation (16). The fact that our two  $k_{10}$  values (calculated using a stirred reactor and an unstirred reactor concept) bracketed the other two recently reported  $k_{10}$  values suggests that the reactor reported herein had incomplete mixing.

The  $\Delta E$  estimate of 1200 calories per mole (5000 J/mole) can be compared with two recent  $\Delta E$  estimates of 600 and 1700 calories per mole (2500 and 7100 J/mole) (refs. 14 and 15, respectively) for this reaction. These referenced values were both obtained by combining measured  $k_{10}$  values at 300° K with some other  $k_{10}$  values obtained by workers using flame studies at the higher temperatures.

The possibility that some of the  $\text{CO}_2$  produced in the  $\text{CO} + \text{OH}$  experiment was due to the action of  $\text{HO}_2$ , the hydroperoxy radical, was eliminated by reacting H with a mixture of  $\text{O}_2 + \text{CO}$  instead of  $\text{NO}_2 + \text{CO}$ . Only a minor amount of  $\text{CO}_2$  was produced for our experimental condition. An approximate rate for the  $\text{CO} + \text{HO}_2$  reaction at room temperature can be obtained by making a crude estimate that the  $+\Delta[\text{CO}]_2$  was only about 1/50 of the  $+\Delta[\text{CO}_2]$  for the  $\text{CO} + \text{OH}$  reaction and that the  $[\text{HO}_2]_t$  in the stirred reactor was about the same as  $[\text{OH}]_t$  in the previous experiments. Then the rate for the  $\text{CO} + \text{HO}_2$  reaction is about  $10^8$  to  $10^9$  cubic centimeters per mole per second.

## Rate Constants for CO + O Reaction

Most of the measurements for this reaction were conducted using a 2055-cubic-centimeter reactor and atomic oxygen derived from a 1.46-percent O<sub>2</sub>-Ar gas mixture. For a temperature range of 400° to 500° K, pressures of 0.75 to 3.5 millimeters (100 to 465 N/m<sup>2</sup>), and amounts of added CO to a mole fraction of 0.15, the loss of atomic oxygen due to reaction with CO was too small to be detected. Traces of CO<sub>2</sub> were observed at the lower limit of detection of the mass spectrometer. It was estimated that the mole fraction of CO<sub>2</sub> produced was about  $\leq 10^{-3}$  at 500° K. Similar results were obtained using atomic oxygen from the N + NO titration technique and from discharged 100-percent O<sub>2</sub>.

The reaction of CO and O is expected to occur primarily by a termolecular process, with a third body required to carry away the heat of reaction. This is contrary to the results of Mahan and Solo (ref. 4) who found the reaction to be apparently bimolecular. In spite of their results, because of the strong theoretical expectation that the reaction is termolecular, the rate constant reported herein has been calculated for a termolecular reaction. For this calculation, the upper limit mole fraction value of  $10^{-3}$  for CO<sub>2</sub> production, the observed oxygen atom mole fraction value of 0.014,  $\Delta t = 1$  second, the added CO mole fraction value of 0.15, and a total pressure of 1 millimeter (133 N/m<sup>2</sup>) were used. The rate constant that resulted was  $4 \times 10^{13}$  cubic centimeters squared per mole squared per second. This value represents an upper limit, because of uncertainty in estimating the background ion current for CO<sub>2</sub>. Other rate constants from the literature are listed for comparison in table II. The bimolecular rate constant of Mahan and Solo (ref. 4) has been recalculated as a termolecular rate constant at a total pressure of 1 millimeter (133 newtons/meter<sup>2</sup>) to allow a direct comparison with the termolecular rate constants

TABLE II. - RATE CONSTANTS FOR CARBON MONOXIDE PLUS OXYGEN  
ATOM REACTIONS

Investigator	Rate constant, k, cm <sup>6</sup> mole <sup>-2</sup> sec <sup>-1</sup>	Temperature, °K
Mahan and Solo <sup>a</sup> , 1962 (ref. 4)	$7 \times 10^{15}$	500
Avramenko and Kolesnikova, 1959 (ref. 3)	$8 \times 10^{16}$	500
Kaufman, 1961 (ref. 24)	$< 5 \times 10^{14}$	300
Clyne and Thrush, 1963 (ref. 25)	$< 3 \times 10^{13}$	300
Groth <sup>b</sup> , 1937 (ref. 26)	$1 \times 10^{13}$	300
Kondrat'ev and Ptichkin, 1961 (ref. 27)	$5 \times 10^{13}$	500
Brabbs and Belles, 1966 (ref. 28)	$2 \times 10^{13}$	3000
This research, 1967	$\leq 4 \times 10^{13}$	500

<sup>a</sup>Reported reaction to be bimolecular.

<sup>b</sup>Relative rate to  $O + O_2 + M \rightarrow O_3 + M$ .



reported by others. Large differences are evident in table II, with rate constants ranging from  $10^{13}$  to  $10^{16}$  cubic centimeters squared per mole squared per second. The general weight of the results favor a low value, less than about  $5 \times 10^{13}$  cubic centimeters squared per mole squared per second.

## Effect of Traces of Hydrogen Compounds on CO + O Reaction

The large differences between different measurements of the CO + O rate constant are outside experimental error. We believe that this might be due to hydrogen compounds in the system, resulting in the presence of OH. Since OH reacts so rapidly with CO, traces of this compound will greatly affect the oxidation rate of CO. Using the rate constants obtained in the present investigation shows that, at  $500^\circ$  K and a pressure of 1 millimeter ( $133 \text{ N/m}^2$ ), the rate of  $\text{CO}_2$  formation by O and OH are equal when the ratio of [O] to [OH] is  $10^5$ . Thus, for experimental conditions, where the atomic oxygen mole fraction is usually about 0.014, a mole fraction of OH larger than  $1 \times 10^{-7}$  could result in most of the  $\text{CO}_2$  being due to the CO + OH reaction. This indicates how difficult it is to make reliable CO + O rate measurements, unless extreme care is used to remove traces of moisture or other hydrogen compounds from the system. Moisture in the oxygen will yield OH when the mixture is passed through the discharge. Hydrogen or hydrocarbons in the CO will yield OH by reaction with atomic oxygen. In either case, it seems possible that none of the rate constants reported for the CO + O oxidation are correct. All observed  $\text{CO}_2$  formation may be due to the presence of traces of OH. In order to gain some experimental evidence of the importance of moisture, the  $\text{O}_2$ -Ar mixture was bubbled through water at room temperature at 1 atmosphere pressure ( $1 \times 10^5 \text{ N/m}^2$ ) before being passed through the microwave discharge. The amount of  $\text{CO}_2$  produced upon reaction with CO in the reactor was doubled for this "wet" mixture. Evidently, OH was present in large enough amounts to influence considerably the reaction rate.

The rate constant reported by Mahan (ref. 4) is about 100 times larger than the observations reported herein, possibly indicating the presence of OH in Mahan's system. Evidence for this can be seen from some of Mahan's experiments in which  $\text{NO}_2$  was introduced into the reactor with CO and O. The addition of  $\text{NO}_2$  increased the  $\text{CO}_2$  production by a factor of about nine. This large effect might be due to the CO + OH reaction, in which the OH was generated by a cyclic process involving the reaction of  $\text{H} + \text{NO}_2 \rightarrow \text{OH} + \text{NO}$ , followed by reaction of OH with CO to form  $\text{CO}_2$  and H. The phenomenon of an increase in  $\text{CO}_2$  production when  $\text{NO}_2$  is added to a CO + O system was also observed recently by Hartek, Reeves, and Thompson (ref. 21). The fact that Mahan observed a bimolecular rate constant might also arise from the dominance by OH of the oxidation process.

## SUMMARY OF RESULTS

Rate measurements were made on the  $\text{CO} + \text{OH}$  and  $\text{CO} + \text{O}$  reactions with the following results:

1. For the  $\text{CO} + \text{OH} \rightarrow \text{CO}_2 + \text{H}$  reaction the rate constant  $k_{10}$  based on  $\text{CO}_2$  production was determined to be  $(4.2 \pm 0.8) \times 10^{11} \exp(-1200 \pm 200/RT)$  cubic centimeters per mole per second. The rate constant value at  $300^\circ \text{K}$  is somewhat lower than two recent measurements of this value possibly because the reactor was not completely mixed. The measured  $\Delta E$  of 1200 calories per mole (5000 J/mole) agrees with the more recent  $\Delta E$  estimates of 0 to 2000 calories per mole (0 to 8333 J/mole) for this reaction.

2. For the  $\text{CO} + \text{O} + \text{M} \rightarrow \text{CO}_2 + \text{M}$  reaction, an upper limit estimate of its rate constant at  $500^\circ \text{K}$  was found to be  $4 \times 10^{13}$  cubic centimeters squared per mole squared per second. This value agrees approximately with some published results, but is much smaller than other published rate constant values. The wide scatter in published rate constant values is believed due to the presence of hydrogen compounds in the various experimenters' reaction systems.

Lewis Research Center,  
National Aeronautics and Space Administration,  
Cleveland, Ohio, June 2, 1967,  
129-01-03-01-22.

## REFERENCES

1. Lewis, Bernard; and von Elbe, Guenther: Combustion, Flames, and Explosions of Gases. Second ed., Academic Press Inc., 1961, p. 75.
2. Reeves, Robert R., Jr.; Harteck, Paul; Thompson, Barbara A.; and Waldron, Roger W.: Photochemical Equilibrium Studies of Carbon Dioxide and their Significance for the Venus Atmosphere. J. Phys. Chem., vol. 70, no. 5, May 1966, pp. 1637-1640.
3. Avramenko, L. I.; and Kolesnikova, R. V.: Kinetics and Mechanism for the Reaction of Oxygen Atoms with Carbon Monoxide. Izvest. Akad. Nauk SSSR, Otdel. Khim. Nauk, vol. 9, 1959, pp. 1562-1570.
4. Mahan, Bruce H.; and Solo, Richard B.: Carbon Monoxide-Oxygen Atom Reaction. J. Chem. Phys., vol. 37, no. 11, Dec. 1, 1962, pp. 2669-2677.
5. Ashmore, P. G.: Elementary Combustion Reactions. Neutral Species. Tenth Symposium (International) on Combustion. The Combustion Institute, Pittsburgh, 1965, pp. 377-386.

6. Baldwin, R. R.; Jackson, D.; Waker, R. W.; and Webster, S. J.: The Use of the Hydrogen-Oxygen Reaction in Evaluating Velocity Constants. Tenth Symposium (International) on Combustion. The Combustion Institute, Pittsburgh, 1965, pp. 423-433.
7. Blundell, R. V.; Cook, W. G. A.; Hoare, D. E.; and Milne, G. S.: Rates of Radical Reactions in Methane Oxidation. Tenth Symposium (International) on Combustion. The Combustion Institute, Pittsburgh, 1965, pp. 445-452.
8. Dixon-Lewis, G.; Sutton, M. M.; and Williams, A.: Some Reactions of Hydrogen Atoms and Simple Radicals at High Temperatures. Tenth Symposium (International) on Combustion. The Combustion Institute, Pittsburgh, 1965, pp. 495-502.
9. Avramenko, L. I.; and Lorentso, R. V.: Reaction of Hydroxyl With the Molecules of Various Gases. II. Reactions of Free Hydroxyl With Hydrogen and Carbon Monoxide. Zhur. Fiz. Khim., vol. 24, 1950, pp. 207-213.
10. Kondrat'ev, V. N.: Reactions of Some Radicals and Their Concentration in Flames. Seventh Symposium (International) on Combustion. Butterworth and Co. (London), 1959, pp. 41-44.
11. Fristrom, R. M.: Radical Concentrations and Reactions in a Methane-Oxygen Flame. Ninth Symposium (International) on Combustion. W. G. Berl, ed., Academic Press, 1963, pp. 560-575.
12. Fenimore, C. P.; and Jones, G. W.: The Reaction of Hydrogen Atoms with Carbon Dioxide at 1200-1350° K. J. Phys. Chem., vol. 62, no. 12, Dec. 1958, pp. 1578-1581.
13. Fenimore, C. P.; and Jones, G. W.: The Decomposition of Ethylene and Ethane in Pre-mixed Hydrocarbon-Oxygen-Hydrogen Flames. Ninth Symposium (International) on Combustion. W. G. Berl, ed., Academic Press, 1963, pp. 597-606.
14. Dixon-Lewis, G.; Wilson, W. E.; and Westenberg, A. A.: Studies of Hydroxyl Radical Kinetics by Quantitative ESR. J. Chem. Phys., vol. 44, no. 8, Apr. 15, 1966, pp. 2877-2884.
15. Herron, John T.: Mass-Spectrometric Study of the Rate of the Reaction  $\text{CO} + \text{OH}$ . J. Chem. Phys., vol. 45, no. 5, Sept. 1, 1966, pp. 1854-1855.
16. Kaufman, F.; and Del Greco, F. P.: Fast Reactions of OH Radical. Ninth Symposium (International) on Combustion. W. G. Berl, ed., Academic Press, 1963, pp. 659-668.
17. Wong, E. L.; and Potter, A. E.: Mass-Spectrometric Investigation of the Reactions of O Atoms with  $\text{H}_2$  and  $\text{NH}_3$ . J. Chem. Phys., vol. 43, no. 9, Nov. 1, 1965, pp. 3371-3382.

18. Phillips, L. F.; and Schiff, H. I.: Mass Spectrometric Studies of Atomic Reactions. III. Reactions of Hydrogen Atoms with Nitrogen Dioxide and with Ozone. J. Chem. Phys., vol. 37, no. 6, Sept. 15, 1962, pp. 1233-1238.
19. Westenberg, A. A.; and de Hass, H.: Quantitative ESR Measurements of Gas-Phase H and OH Concentrations in the H-NO<sub>2</sub> Reactions. J. Chem. Phys., vol. 43, no. 5, Sept. 1, 1965, pp. 1550-1556.
20. Westenberg, A. A.; and Fristrom, R. M.: Methane-Oxygen Flame Structure. IV. Chemical Kinetic Considerations. J. Phys. Chem., vol. 65, no. 4, Apr. 1961, pp. 591-601.
21. Harteck, P.; Reeves, R. R., Jr.; and Thompson, B. A.: Chemistry of Planetary Atmosphere and Comets. Tech. Status Rep. No. 7, Aug. 1, 1965 - Feb. 28, 1966, Rensselaer Poly. Inst., 1966. (NASA Grant NsG-261.)
22. Dixon-Lewis, G.; Sutton, M. M.; and Williams, Alan: The Reaction of Hydrogen Atoms with Nitrous Oxide. J. Chem. Soc., Sept. 1965, pp. 5724-5729.
23. Jost, W.; Schecker, H. G. G.; and Wagner, H. G. G.: Measurement of the Rate of the Regulation of the Water Gas Equilibrium. Z. Physik. Chem., vol. 45, no. 1-2, 1965, pp. 47-57.
24. Kaufman, Frederick: Reactions of Oxygen Atoms. Progress in Reaction Kinetics. Vol. I., G. Porter, ed., Pergamon Press, 1961, pp. 3-39.
25. Clyne, M. A. A.; and Thrush, B. A.: The Kinetics of the Carbon Monoxide Flame Bands. Ninth Symposium (International) on Combustion. W. G. Berl, ed., Academic Press, 1963, pp. 177-183.
26. Groth, W.: Photochemical Experiments in the Schumann Ultraviolet. IV. The Photochemical Oxidation of Carbon Monoxide and Carbon Monoxide as a Sensitizer. Z. Physik. Chem., vol. B37, 1937, pp. 315-322.
27. Kondrat'ev, V. N.; and Ptichkin, I. I.: Gas-Phase Interaction Between Carbon Monoxide and Ozonized Oxygen. Kinetika i Kataliz, vol. 2, July-Aug. 1961, pp. 449-452.
28. Brabbs, Theodore A.; and Belles, Frank E.: Recombination of Carbon Monoxide and Atomic Oxygen at High Temperatures. NASA TN D-3744, 1966.

*"The aeronautical and space activities of the United States shall be conducted so as to contribute . . . to the expansion of human knowledge of phenomena in the atmosphere and space. The Administration shall provide for the widest practicable and appropriate dissemination of information concerning its activities and the results thereof."*

—NATIONAL AERONAUTICS AND SPACE ACT OF 1958

## NASA SCIENTIFIC AND TECHNICAL PUBLICATIONS

**TECHNICAL REPORTS:** Scientific and technical information considered important, complete, and a lasting contribution to existing knowledge.

**TECHNICAL NOTES:** Information less broad in scope but nevertheless of importance as a contribution to existing knowledge.

**TECHNICAL MEMORANDUMS:** Information receiving limited distribution because of preliminary data, security classification, or other reasons.

**CONTRACTOR REPORTS:** Scientific and technical information generated under a NASA contract or grant and considered an important contribution to existing knowledge.

**TECHNICAL TRANSLATIONS:** Information published in a foreign language considered to merit NASA distribution in English.

**SPECIAL PUBLICATIONS:** Information derived from or of value to NASA activities. Publications include conference proceedings, monographs, data compilations, handbooks, sourcebooks, and special bibliographies.

**TECHNOLOGY UTILIZATION PUBLICATIONS:** Information on technology used by NASA that may be of particular interest in commercial and other non-aerospace applications. Publications include Tech Briefs, Technology Utilization Reports and Notes, and Technology Surveys.

*Details on the availability of these publications may be obtained from:*

SCIENTIFIC AND TECHNICAL INFORMATION DIVISION  
NATIONAL AERONAUTICS AND SPACE ADMINISTRATION  
Washington, D.C. 20546